


André Lessa BSM Theory (UFABC – São Paulo)


SPACE Workshop

October 24th, 2025

Research Projects




- The tool was developed by theorists and experimentalists
- It has an active group of collaborators
- It is in constant development




- Largest database of BSM searches:
 - 61 ATLAS results
 - 64 CMS results

- The tool was developed by theorists and experimentalists
- It has an active group of collaborators
- It is in constant development



- Largest database of BSM searches:
 - 61 ATLAS results
 - 64 CMS results





- Recent developments
 - Full statistical models (Pyhf)
 - Global analysis combination
 - Arbitrary models → graph theory




- Recent developments
 - Full statistical models (Pyhf)
 - Global analysis combination
 - Arbitrary models → graph theory

• Future plans:


- Inclusion of Run 3 results
- More LLP (non-SUSY) results
- Impact of analysis combination
- ...

• The large database of ATLAS and CMS results allow for a meta-analysis:



- Currently the largest excess are at $\sim 3-4\sigma$
- These are expected due to the huge number of searches! (~1200 bins)


• The large database of ATLAS and CMS results allow for a meta-analysis:



- Currently the largest excess are at $\sim 3-4\sigma$
- These are expected due to the huge number of searches! (~1200 bins)

- However, what if:
 - Signal is dispersed over many final states (not easily visible by individual searches)?
 - The Next SM (NSM) does not match any of the UV models considered so far?
 - → How to infer the NSM from data?


- We have built a prototype algorithm based on a MCMC for a dimensional space of varying dimensions.
- The algorithm looks for dispersed signals which can be simultaneously explained by minimal number of new particles.



S. Kraml, AL, W. Waltenberger, *JHEP 03 (2021) 207*

Feedback for building new models

- We have built a prototype algorithm based on a MCMC for a dimensional space of varying dimensions.
- The algorithm looks for dispersed signals which can be simultaneously explained by minimal number of new particles.



S. Kraml, AL, W. Waltenberger, *JHEP 03 (2021) 207*

Feedback for building new models

Future plans:


- New version under development
- Reversible-Jump MCMC
- Better statistical interpretation
- Experimental bias
- How to automatically build a Lagragian (FeynRules → UFO)?
-

• A common approach is to make use of EFTs to look for NP on SM measurements. However...

- A common approach is to make use of EFTs to look for NP on SM measurements. However...
 - Is the EFT assumption valid/consistent?

- A common approach is to make use of EFTs to look for NP on SM measurements. However...
 - Is the EFT assumption valid/consistent?
 - Are the EFT constraints valid even if the EFT assumption is violated?
 - If not, are they too conservative/too aggressive?
 - What are we missing by looking only for the on-shell or EFT regimes?

- A common approach is to make use of EFTs to look for NP on SM measurements. However...
 - Is the EFT assumption valid/consistent?
 - Are the EFT constraints valid even if the EFT assumption is violated?
 - If not, are they too conservative/too aggressive?
 - What are we missing by looking only for the on-shell or EFT regimes?
 - What happens in the transition regime? $\left(\Lambda_{
 m BSM}\gtrsim\sqrt{\hat{s}}
 ight)$



$$\mathcal{L}_{BSM} = |D_{\mu}\phi_T|^2 - y_{\mathrm{DM}}\phi_T^{\dagger} \bar{\chi} t_R + ...$$
 On-shell


$$q \qquad t$$

$$\bar{q} \qquad \bar{t}$$

$$C_{\text{SMEFT}} = \sum_{i} \frac{c_i^{(6)}}{\Lambda^2} O_i^{(6)} + \dots$$

$$EFT$$

- A common approach is to make use of EFTs to look for NP on SM measurements. However...
 - Is the EFT assumption valid/consistent?
 - Are the EFT constraints valid even if the EFT assumption is violated?
 - If not, are they too conservative/too aggressive?
 - What are we missing by looking only for the on-shell or EFT regimes?
 - What happens in the transition regime? $\left(\Lambda_{
 m BSM}\gtrsim\sqrt{\hat{s}}
 ight)$



$$\mathcal{L}_{BSM} = |D_{\mu}\phi_T|^2 - y_{\mathrm{DM}}\phi_T^{\dagger}ar{\chi}t_R + ...$$

On-shell

$$\bar{q}$$
 ϕ_T
 χ
 $\bar{\chi}$

 $2m_T \simeq \sqrt{\hat{s}}$

$$\mathcal{L}_{BSM} = |D_{\mu}\phi_{T}|^{2} - y_{\mathrm{DM}}\phi_{T}^{\dagger}\bar{\chi}t_{R} + \dots \qquad \mathcal{L}_{\mathrm{FF}} = \sum_{i} \mathcal{F}_{i}(q^{2},\Lambda^{2},\dots)O_{i}^{(6)} + \dots \qquad \mathcal{L}_{\mathrm{SMEFT}} = \sum_{i} \mathcal{F}_{i$$

$$\frac{q}{\bar{q}} \qquad t$$

$$\bar{t}$$


$$\text{MEFT} = \sum_{i} \frac{c_i^{(6)}}{\Lambda^2} O_i^{(6)} + ...$$

 $2m_T \gg \sqrt{\hat{s}}$

Beyond EFTs

• Form factors encode the full kinematical dependence at any energy:

$$\mathcal{F}_i(q^2,\Lambda^2,\ldots) \sim c_i^0 \frac{1}{\Lambda^2} + c_i^1 \frac{q^2}{\Lambda^4} + \ldots$$
 New features not captured by EFTs!




- New features in SM distributions:
 - Broad resonance
 - Deficit of events at higher bins (negative interference)

Beyond EFTs

Form factors encode the full kinematical dependence at any energy:

$$\mathcal{F}_i(q^2,\Lambda^2,\ldots) \sim c_i^0 \frac{1}{\Lambda^2} + c_i^1 \frac{q^2}{\Lambda^4} + \ldots$$
 New features not captured by EFTs




- captured by EFTs!
- New features in SM distributions:
 - Broad resonance
 - Deficit of events at higher bins (negative interference)

Future plans

- New search strategies (Anomaly detection?)
- Combining EFT (low energy) and form factors (high energy) constraints
- Model dependence → generic features?
- Other observables?

New Signatures - LLPs




45

40

35

15

• There are several "minimal" scenarios which lead to LLP signatures.



 $m_X ({
m GeV})$

DD Excluded

 $\Omega_{DM}h^2 = 0.12$

WIMP Region

CDFO Region

J. Heisig, AL, L. Magno, PRD 110 (2024) 1

New Signatures - LLPs

- Many of these scenarios are not yet fully explored by searches.
- In the last years searches for LLPs have increased considerably at the LHC.
- The collaboration between the experimental and theory communities has been key for developing the program.

LLP Working Group (CERN-LPCC):

- Theory: André Lessa
- ALICE: Mesut Arslandok
- ATLAS: Dominique Trischuk
- CMS: Alberto Escalante del Valle
- FASER: Dave Casper
- LHCb: Gaia Lanfranchi and Andrii Usachov
- MoEDAL: James Pinfold
- SND@LHC: Cristovao Vilela

New Signatures - LLPs

- Many of these scenarios are not yet fully explored by searches.
- In the last years searches for LLPs have increased considerably at the LHC.
- The collaboration between the experimental and theory communities has been key for developing the program.

LLP Working Group (CERN-LPCC):


- Theory: André Lessa
- ALICE: Mesut Arslandok
- ATLAS: Dominique Trischuk
- CMS: Alberto Escalante del Valle
- FASER: Dave Casper
- LHCb: Gaia Lanfranchi and Andrii Usachov
- MoEDAL: James Pinfold
- SND@LHC: Cristovao Vilela


Future plans

- New trigger strategies
- Coordination of benchmarks for searches
- Complementarity between prompt and LLP searches
- New signatures/gaps in coverage


Backup

• Current run:





Analysis Name	Type	Dataset	Topos	Observed	Expected	Approx σ
CMS-EXO-20-004	comb -		TChiISR, TChiZISRqq	2e+02 fb	1e+02 fb	4.6 σ
ATLAS-EXOT-2018-06	em I	EM10	TChiISR, TChiZISRqq	413	359.00 +/- 10.00	2.5 σ
CMS-SUS-20-004	comb -		TChiHH	0.3 fb	0.2 fb	2.2 σ
CMS-SUS-13-012	em 3	3NJet6_1000HT12	TChiZZ	32	22.80 +/- 5.20	1.3 σ



EFT vs Direct Searches

- What happens when we consider complex scenarios? (MSSM)
- Are the EFT constraints valid/competitive?

$$\mathcal{L}_{ ext{MSSM}}(m_{\tilde{q}},...) \Rightarrow \mathcal{L}_{ ext{SMEFT}} = \mathcal{L}_{ ext{SM}} + \sum_{i} rac{C_{i}^{(6)}(m_{\tilde{q}},...)}{\Lambda^{2}} O_{i}^{(6)} + \mathcal{O}\left(\Lambda^{-4}\right)$$

MSSM

(4-component spinors, diagonal Higgs basis)



Functional methods (CDE)

SMEFT


(Warsaw basis)

S. Kraml, AL, S. Prakash, F. Wilsch, arXiv:2506.05201

```
C_{HB} \rightarrow \hbar \left( \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{3,0} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{2} g_{1}^{2} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{2} g_{1}^{2} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{2} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{2} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{2} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{2} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{2} LF_{4,-1} \left[ m_{\tilde{d}}^{p} \right] - \frac{1}{36} \sum_{p} c_{2\beta} g_{1}^{2} LF_{4,-1} \left[ m_{\tilde{d}}^{p} 
                                                              \frac{1}{6} \; {g_{1}}^{2} \; {c_{\beta}}^{2} \; \overline{y_{d}}^{pr} \; {y_{d}}^{pr} \; \mathsf{LF_{3,0}} \left[ \, m_{\tilde{d}}^{\; r} \, \right] \; + \\ \frac{1}{6} \; {g_{1}}^{2} \; {c_{\beta}}^{2} \; \overline{y_{d}}^{pr} \; {y_{d}}^{pr} \; \mathsf{LF_{4,-1}} \left[ \, m_{\tilde{d}}^{\; r} \, \right] \; + \\ \frac{1}{6} \; {g_{1}}^{2} \; {c_{\beta}}^{2} \; \overline{y_{d}}^{pr} \; {y_{d}}^{pr} \; \mathsf{LF_{4,-1}} \left[ \, m_{\tilde{d}}^{\; r} \, \right] \; + \\ \frac{1}{6} \; {g_{1}}^{2} \; {c_{\beta}}^{2} \; \overline{y_{d}}^{pr} \; {y_{d}}^{pr} \; {b_{1}}^{pr} \; {b_{2}}^{pr} \; {b
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            \frac{1}{4} g_1^2 \left( c_\beta \overline{a_e}^{pr} - s_\beta \widetilde{\mu} \overline{y_e}^{pr} \right) \left( c_\beta a_e^{pr} - s_\beta \widetilde{\mu} y_e^{pr} \right) LF_{5,1,-2} \left[ m_{\tilde{1}}^{p}, m_{\tilde{e}}^{r} \right] +
                                                            \frac{1}{4} \sum_{\mathbf{p}} c_{2\,\beta} \, g_{1}^{\ 4} \, \mathsf{LF}_{3,\,\theta} \big[ \, \mathsf{m}_{\check{e}}^{\ p} \, \big] \, - \, \frac{1}{4} \sum_{\mathbf{p}} c_{2\,\beta} \, g_{1}^{\ 4} \, \mathsf{LF}_{4,\,-1} \big[ \, \mathsf{m}_{\check{e}}^{\ p} \, \big] \, - \, \frac{1}{2} \, g_{1}^{\ 2} \, c_{\beta}^{\ 2} \, \overline{\mathsf{y}_{e}}^{\, pr} \, \mathsf{y}_{e}^{\, pr} \, \mathsf{LF}_{3,\,\theta} \big[ \, \mathsf{m}_{\check{e}}^{\, r} \, \big] \, + \, \frac{1}{2} \, \mathsf{g}_{1}^{\ 2} \,
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            \frac{1}{3} g_1^2 \left( c_\beta \overline{a_d}^{pr} - s_\beta \widetilde{\mu} \overline{y_d}^{pr} \right) \left( c_\beta a_d^{pr} - s_\beta \widetilde{\mu} y_d^{pr} \right) LF_{3,1,0} \left[ m_{\widetilde{u}}^p, m_{\widetilde{d}}^r \right] +
                                                                 \frac{1}{2} g_1^2 c_{\beta}^2 \overline{y_e}^{pr} y_e^{pr} LF_{4,-1} [m_{\tilde{e}}^r] - \frac{1}{16} g_1^2 (2 q)
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              (c_{\beta} a_d^{pr} - s_{\beta} \tilde{\mu} y_d^{pr}) LF_{3,2,-1}[m_{\tilde{q}}^p, m_{\tilde{d}}^r] -
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 OperatorToC++
                                                               \frac{1}{16} \; \left( 2 \; g_1^{\; 2} \; c_\beta^{\; 2} \; \overline{y_e}^{pr} \; y_e^{pr} \; + \sum_p \, c_{2 \, \beta} \; g_1^{\; 4} \right) \; \mathsf{LF_4},_{-1}
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               \left(c_{\beta} a_{d}^{pr} - s_{\beta} \widetilde{\mu} y_{d}^{pr}\right) LF_{4,1,-1}\left[m_{\widetilde{q}}^{p}, m_{\widetilde{d}}^{r}\right] +
                                                               \frac{1}{144} \ g_1^2 \ \left( - 6 \ c_{eta}^{\ 2} \ \overline{y_d}^{pr} \ y_d^{pr} - 6 \ s_{eta}^{\ 2} \ \overline{y_u}^{pr} \ y_u^{pr} + \right.
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            \left(c_{\beta} a_{d}^{pr} - s_{\beta} \widetilde{\mu} y_{d}^{pr}\right) LF_{5,1,-2}\left[m_{\tilde{a}}^{p}, m_{\tilde{d}}^{r}\right] -
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     allows for the numerical
                                                              \frac{1}{144} g_1^2 \left( 6 c_\beta^2 \overline{y_d}^{pr} y_d^{pr} + 6 s_\beta^2 \overline{y_u}^{pr} y_u^{pr} - \sum \right)
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            (s_{\beta} a_{u}^{pr} - \tilde{\mu} c_{\beta} y_{u}^{pr}) LF_{2,2,0}[m_{\tilde{a}}^{p}, m_{\tilde{u}}^{r}] -
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   evaluation of coefficients given
                                                                 \frac{2}{9} \sum_{p} c_{2\beta} g_{1}^{4} LF_{3,0} [m_{\tilde{u}}^{p}] + \frac{2}{9} \sum_{p} c_{2\beta} g_{1}^{4} LF_{4,}]
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               (s_{\beta} a_{u}^{pr} - \tilde{\mu} c_{\beta} y_{u}^{pr}) LF_{3,1,0}[m_{\tilde{q}}^{p}, m_{\tilde{u}}^{r}] +
                                                                              g_1^2 s_\beta^2 \overline{y_u}^{pr} y_u^{pr} LF_{4,-1} [m_{\tilde{u}}^r] + \frac{1}{64} (g_1^4)
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               (s_{\beta} a_{u}^{pr} - \widetilde{\mu} c_{\beta} y_{u}^{pr}) LF_{3,2,-1}[m_{\widetilde{a}}^{p}, m_{\widetilde{u}}^{r}] +
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   the input parameters
                                                                 \frac{1}{64} g_1^2 (g_1^2 (1 + 3 c_{4\beta}) + 3 g_2^2 (-1 + c_{4\beta}))
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 (s_{\beta} a_{u}^{pr} - \tilde{\mu} c_{\beta} y_{u}^{pr}) LF_{4,1,-1}[m_{\tilde{0}}^{p}, m_{\tilde{\mu}}^{r}] -
                                                                               g_1^2 \left( c_\beta \overline{a_d}^{pr} - s_\beta \widetilde{\mu} \overline{y_d}^{pr} \right) \left( c_\beta a_d^{pr} - s_\beta \widetilde{\mu} y_d \right)
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       (s_{\beta} a_{u}^{pr} - \widetilde{\mu} c_{\beta} y_{u}^{pr}) LF_{3,1,0}[m_{\widetilde{u}}^{r}, m_{\widetilde{a}}^{p}] +
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       \frac{1}{24} g_1^2 \left( s_\beta \overline{a_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) \left( s_\beta a_u^{pr} - \widetilde{\mu} c_\beta y_u^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^r, m_{\widetilde{q}}^{-p} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{a_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{-r}, m_{\widetilde{q}}^{-p} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{a_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{-r}, m_{\widetilde{q}}^{-p} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{a_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{-r}, m_{\widetilde{q}}^{-p} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{a_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{-r}, m_{\widetilde{q}}^{-p} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{a_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{-r}, m_{\widetilde{q}}^{-p} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{a_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{-r}, m_{\widetilde{q}}^{-p} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{-r}, m_{\widetilde{q}}^{-p} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{-r}, m_{\widetilde{q}}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,-1} \left[ m_{\widetilde{u}}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right] - \frac{1}{24} g_1^2 \left( s_\beta \overline{y_u}^{pr} - \widetilde{\mu} c_\beta \overline{y_u}^{pr} \right) LF_{3,2,
```

EFT vs Direct Searches






Future plans

- EFT constraints vs direct searches
- Higgs sector (2HDM) vs EFT
- Multi-scale MSSM


Main collaborators:


- S. Kraml (LPSC-Grenoble)
- S. Prakash (IFIC)
- F. Wilsch (RWTH-Aachen)

• What is the real impact of LHC data on BSM models?




- What is the real impact of LHC data on BSM models?
 - → Not an easy question!






• SModelS → software tool for collecting experimental results and allowing for a quick assessment of constraints to BSM scenarios




• SModelS → software tool for collecting experimental results and allowing for a quick assessment of constraints to BSM scenarios



SModelS: Basics

 Tool for reintrepretation of LHC results based on rescaling of simplified model (SMS) results:



• Pros:

- Many SMS results have been produced by the experimental collaborations
- No need for recasting or event simulation
- Sometimes the only alternative for complex searches
- Very fast!

Cons:


- The rescaling extrapolates/interpolates signal efficiencies
- Results are limited by the available simplified models
- Results tend to be conservative and underestimate the experimental sensitivity

SModelS: Basics

- SModelS performs 3 main tasks:
 - 1. UV model → sum of SMS
 - 2. Find the matching results in the database
 - 3. Compute constraints (limit setting)


Main approximation:

SMS efficiencies are assumed to be driven by the properties of on-shell particles a masses, widths and quantum numbers



SModelS Graphs

• Graph Representation:





• Text Representation:

SModelS v3: Database

• Non-Z₂ results:


ID	Signature	Luminosity	SMS Topology	Type		
Run 2 - 13 TeV						
ATLAS-EXOT-2019-03	Dijet resonance	$139 \; {\rm fb}^{-1}$	$pp o Z' o jj, bar{b}$	UL		
ATLAS-EXOT-2018-48	$t\bar{t}$ resonance	$139 \; {\rm fb^{-1}}$	pp o Z' o t ar t	UL		
CMS-EXO-19-012	Dijet resonance	$137 \; {\rm fb^{-1}}$	$pp o Z' o jj, b\bar{b}$	UL		
CMS-EXO-20-008	<i>b</i> -jet resonance	$138 \; {\rm fb^{-1}}$	$pp o Z' o b \overline{b}$	UL		
CMS-EXO-20-004	Monojet	$137 \; {\rm fb^{-1}}$	$pp \to Z', S \to \chi \chi$	EM		
ATLAS-EXOT-2018-06	Monojet	$139 \; {\rm fb^{-1}}$	$pp \to Z' \to \chi \chi$	UL		
ATLAS-SUSY-2018-22	Multi-jet plus	$139 \; {\rm fb^{-1}}$	$pp \to Z' \to \chi \chi$	EM		
ATLAS-SUSY-2018-13	Displaced jets	$139 \; {\rm fb^{-1}}$	$pp \to \tilde{\chi}\tilde{\chi} \to jjj, jjj; \dots$	EM		
$\operatorname{Run} 1$ - 8 TeV						
CMS-EXO-16-057	b-jet resonance	$19.7 \; {\rm fb^{-1}}$	$pp o Z' o b \overline{b}$	UL		
CMS-EXO-12-059	Dijet resonance	19.7 fb^{-1}	pp o Z' o jj	UL		
ATLAS-EXOT-2013-11	Dijet resonance	$20.3 \; {\rm fb^{-1}}$	pp o Z' o jj	UL		







Physics Study: 2MDM

Two mediator DM






Statistical Combination of ATLAS + CMS



- We have built a prototype algorithm based on a MCMC for a dimensional space of varying dimensions.
- The algorithm looks for dispersed signals which can be simultaneously explained by minimal number of new particles.


- We have built a prototype algorithm based on a MCMC for a dimensional space of varying dimensions.
- The algorithm looks for dispersed signals which can be simultaneously explained by minimal number of new particles.




S. Kraml, AL, W. Waltenberger, *JHEP 03 (2021) 207*

Feedback for building new models

- We have built a prototype algorithm based on a MCMC for a dimensional space of varying dimensions.
- The algorithm looks for dispersed signals which can be simultaneously explained by minimal number of new particles.



S. Kraml, AL, W. Waltenberger, JHEP 03 (2021) 207




Feedback for building new models

- With the increase of data (HL-LHC)
 - → SM observables measured at higher precision

- With the increase of data (HL-LHC)
 - → SM observables measured at higher precision
- A common approach is to make use of EFTs to look for NP on SM measurements

- With the increase of data (HL-LHC)
 - → SM observables measured at higher precision
- A common approach is to make use of EFTs to look for NP on SM measurements



Beyont EFTs

Form factors:

$$\mathcal{F}^{\mu} = i \frac{T^{A}}{32\pi^{4}} \left[2 \left(p_{2} p_{1}^{\mu} + p_{1} p_{2}^{\mu} \right) - s \gamma^{\mu} \right] P_{R} \mathcal{F}(s, m_{T}^{2}) \quad (m_{\chi} \to m_{T}, m_{t} \to 0),$$

where:

$$\mathcal{F}(s, m_T^2) \equiv \frac{1}{s^2} \left[m_T^2 L^2(s, m_T^2) + \sqrt{s \left(s - 4m_T^2 \right)} L(s, m_T^2) + 3s \right] \text{ and }$$

$$L(s, m_T^2) \equiv \log \left[\left(\sqrt{s \left(s - 4m_T^2 \right)} + 2m_T^2 - s \right) / (2m_T^2) \right]$$

